

Sympathetic Epidural (Interlaminar) Block

	-	
What is it?	An epidural injection, or interlaminar block, is an injection of a local anesthetic that is placed in your spine. A fluoroscope (x-ray machine) assists the physician in locating the nerve roots thought to be inflamed and contributing to your pain. It can be done in the neck, mid back, low back, or near the tail bone.	
Why is it done?	This injection can be helpful to patients with pain in the neck, mid or low back, arm or leg. Examples of extremity pain include reflex sympathetic dystrophy (RSD), also known as complex regional pain syndrome (abnormal response of a nerve in the arm or leg). It is also helpful for patients who have been diagnosed with shingles (spinal nerve inflammation.)	
How is it done?	When you are in the procedure room, you will be asked to lie on your stomach on a cushioned x-ray table. If you are receiving conscious sedation, an intravenous catheter (tube) will be inserted into your arm. If you will be receiving local anesthesia, a small needle is used to inject a local anesthetic into your skin to number the area where the anesthetic medication will be injected. You may notice stinging for a few seconds. Next, a slightly larger needle is inserted. Dye is then injected so that fluoroscopy (x-ray machine) can be used to confirm that the needle tip is located in the epidural space (the donut shaped space that surrounds the spinal canal which includes the nerve roots that begin in the spinal canal itself). A solution of local anesthetic is then injected. The needle is removed and a small band aid is applied.	
Is there any preparation?	You may receive conscious sedation. Conscious sedation is medication given through an intravenous (IV) catheter (tube) in your arm. The medication will help you relax, but it will not put you to sleep. If you opt to have conscious sedation, you MUST NOT EAT OR DRINK for 4 hours before your procedure. You may, however, take your medication with a small sip of water. Further, you must have someone available to drive you home after the procedure.	
What to wear?	Please wear loose, comfortable clothing and leave all jewelry and other valuables at home.	
How long does it take?	We ask that you arrive 30 - 45 minutes before the scheduled time of your procedure. The procedure lasts about 15 - 30 minutes, and you will be in the recovery area about 15 - 20 minutes.	
i		
Risks of this procedure include, but are	* Pain in the area where the needle(s) was/were inserted, which can last for two to three days. This can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol.	
procedure	* Pain in the area where the needle(s) was/were inserted, which can last for two to three days. This can be treated by using ice and mild analgesics (pain medication)	
procedure include, but are	 Pain in the area where the needle(s) was/were inserted, which can last for two to three days. This can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face and throat, respiratory difficulties and shock. Medication 	
procedure include, but are	 Pain in the area where the needle(s) was/were inserted, which can last for two to three days. This can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. A reaction to the local anesthetic or conscious sedation, including but not limited to: allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. Temporary numbness or weakness in one or both legs. This is normal. You should have assistance with walking and should not drive for 24 hours. 	
procedure include, but are	 Pain in the area where the needle(s) was/were inserted, which can last for two to three days. This can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. A reaction to the local anesthetic or conscious sedation, including but not limited to: allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. Temporary numbness or weakness in one or both legs. This is normal. You should 	

09/12/2015,09/27/2008rdsteachingsheetREVIEWED 3/2016

<u>Instructions Regarding Your Procedure</u> <u>Sympathetic Epidural (Interlaminar) Block</u>

Will I receive a pre-procedure phone call?	A nurse from our office will call you 24-48 hours prior to your procedure. She will ask you a few important medical questions, <u>AND</u> she will confirm your appointment. We must have <u>a verbal</u> <u>confirmation from you 24-48 hours prior to your procedure</u> . Failure to return our call could result in your procedure being rescheduled. The number to call us back is: (315) 703-3480
Where can I obtain more procedure information? Can I drive home from the	Please visit our website www.nyspineandwellness.com . It contains demonstrations of some of the procedures we offer. We also have other helpful information.
procedure?	You MUST have a ride home, and your driver should remain on site. You must not drive or operate heavy machinery for 24 hours.
Can I eat Breakfast? *the box with a check applies to you	☐ Yes, you can eat a light breakfast or light lunch
	☐ No, <u>DO NOT EAT OR DRINK</u> for <u>4</u> hours before your procedure. However, you may take your medicine but only with a <u>small sip of water</u> .
Should I stop taking aspirin?	If you take ASPIRIN or products containing ASPIRIN in doses greater than 325 mg/per day, you must stop taking it 7 days prior to your procedure (examples of ASPIRIN-containing products include but are not limited to: EXCEDRIN, FIORINAL, AND ALKA SELTZER, etc).
Can I take the aspirin	
prescribed for <u>cardiac</u> and or	☐ Yes, however you need to limit the daily dose to 81 mgs per day for the 7 days prior to the
stroke prevention?	day of the procedure.
Can I take my blood thinner?	☐ Our provider has advised you on the number of days to hold the blood thinner you take.
*the box with a check applies to you	Our provider will have to contact the Doctor who has prescribed the blood thinner for you to learn how long it is safe for you to be off the medicine prior to your procedure. One of
	our nurses will call you when we have obtained that information.
Can I take my other	Yes, except for blood thinners .
medications? *the box with a check applies to you	Yes, only with a small sip of water, (you must not take your blood thinner)
What time is my procedure?	You will receive a phone call one or two business days prior to your procedure to remind you
	of the time you should arrive for your procedure (usually 30-40 minutes prior to the scheduled
	time) and the location. PLEASE BE ON TIME.
	*some patients may receive a call to confirm insurance information.
How long will it take?	We ask that you plan $\frac{1}{2}$ -3 hours from the time you arrive at the center to the time you are
	discharged. Please understand that circumstances sometimes arise that are out of our control,
	which cause a longer delay. Please plan accordingly.
What do I bring?	You <i>must</i> bring the following <u>each time</u> you have a procedure: 1. Your current insurance information
	2. A written list of all your current medications including the strength and frequency you take
	them. Please include over the counter medication, vitamins and herbal supplements.
What shall I wear?	Please wear comfortable loose fitting clothing.
Can I wear jewelry?	No. Please leave all jewelry and other valuables at home. We are unable to store them safely.
, ,-	Any body- piercing jewelry must be completely removed, as this may interfere with the x-ray machine.
What do I do if I am ill or	Please call the office 24 hours in advance if you are unable to keep your appointment.
have to cancel?	(315) 552-6700. We do charge patients for no shows.