

AC JOINT (Acromioclavicular Joint) Injection with fluoroscopy

<i>What is it?</i>	This is a small joint that connects the acromion (part of the shoulder) to the end of your clavicle. If this joint experiences injury, arthritis or mechanical stress one may experience knee pain.
<i>Why is it done?</i>	The procedure is done to relieve pain that origins (begins) in the AC Joint or shoulder. It can help relieve the pain, as well as help diagnose the cause of pain.
<i>How is it done?</i>	When you are in the procedure room, you will be asked to lie on your back on a cushioned x-ray table. A fluoroscope (x-ray machine) assists the provider in visualizing the AC Joint under x-ray. A small needle is placed in the joint and a small amount of solution of local anesthetic (numbing medication) and a cortisone derivative (anti-inflammatory medication) is then injected into the joint. A small band aid is applied after procedure is complete and may be removed at your discretion.
<i>What to wear?</i>	Please wear loose, comfortable clothing and leave all jewelry and other valuables at home.
<i>How long will it take?</i>	Please arrive 30 minutes before your appointment time. The procedure lasts about 15.
<i>Can I drive home from the procedure?</i>	You, can drive yourself to and from the procedure unless medication is taken that can impair your driving ability.
<i>Can I eat before?</i>	Yes we encourage you to eat a normal breakfast and or lunch.
<i>Should I stop taking aspirin?</i>	No
<i>Can I take my medications?</i>	Yes
<i>What are the possible side effects of steroid medication?</i>	Administration of steroid medication can cause side effects. They include, but not limited to: hyperglycemia (increased blood sugar), sweats, hot/cold flashes, flushing of the face, hypertension (increased blood pressure), osteoporosis, insomnia, mood swings and Cushing's Syndrome.
<i>Risks of this procedure include, but are not limited to:</i>	<ul style="list-style-type: none"> * Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. * <i>Bleeding, infection and/or bruising in the injection area. Increased pain</i> * A reaction to the local anesthetic, including but not limited to: allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest requiring transport to the emergency room. . * Nerve damage, paralysis, stroke and even death.

03/15/2014rdsteachingsheetREVIEWED 3/2016

PLEASE SEE THE BACK FOR YOUR INSTRUCTIONS

Instructions Regarding Your Procedure

AC JOINT (Acromioclavicular Joint) Injection with fluoroscopy

Will I receive a pre-procedure phone call?	A nurse from our office will call you 24-48 hours prior to your procedure. She will ask you a few important medical questions, AND she will confirm your appointment. We must have a verbal confirmation from you 24-48 hours prior to your procedure. Failure to return our call could result in your procedure being rescheduled. The number to call us back is: (315) 703-3480
Where can I obtain more procedure information?	Please visit our website www.nyspineandwellness.com . It contains demonstrations of some of the procedures we offer. We also have other helpful information.
Can I drive home from the procedure? <i>*the box with a check applies to you</i>	<input type="checkbox"/> You MUST have a ride home, and your driver should remain on site. You must not drive or operate heavy machinery for 24 hours. <input type="checkbox"/> You can drive yourself home.
Can I eat Breakfast?	Yes, you can eat a light breakfast or light lunch.
Should I stop taking aspirin?	If you take ASPIRIN or products containing ASPIRIN in doses greater than 325 mg/per day, you must stop taking it 7 days prior to your procedure (examples of ASPIRIN-containing products include but are not limited to: EXCEDRIN, FIORINAL, AND ALKA SELTZER, etc).
Can I take the aspirin prescribed for <u>cardiac</u> and or <u>stroke</u> prevention?	Yes, you can continue to take your aspirin for cardiac or stroke prevention, not to exceed 325mg per day.
Can I take my blood thinner?	Yes.
Can I take my other medications?	Yes, take your medications as you normally do.
What time is my procedure?	You will receive a phone call one or two business days prior to your procedure to remind you of the time you should arrive for your procedure (usually 30-40 minutes prior to the scheduled time) and the location. PLEASE BE ON TIME. *some patients may receive a call to confirm insurance information.
How long will it take?	15 minutes.
What do I bring?	You must bring the following <u>each time</u> you have a procedure: 1. Your current insurance information 2. A written list of all your current medications including the strength and frequency you take them. Please include over the counter medication, vitamins and herbal supplements.
What shall I wear?	Please wear comfortable loose fitting clothing.
Can I wear jewelry?	No. Please leave all jewelry and other valuables at home. We are unable to store them safely. Any body- piercing jewelry must be completely removed, as this may interfere with the x-ray machine.
What do I do if I am ill or have to cancel?	Please call the office 24 hours in advance if you are unable to keep your appointment. (315) 552-6700. We do charge patients for no shows.