



Ganglion Impar nerve Block with fluoroscopy

What is it?	The ganglion impar is a structure located at the level of the coccyx.
Why is it done?	A ganglion impar block is a procedure used to reduce some of the symptoms of chronic pelvic or rectal pain by blocking nerve impulses.
How is it done?	Once in the procedure room, our nursing staff will position you on your stomach on a narrow table. Your back will be cleaned with iodine and steriley draped. The doctor will numb your skin with a small needle, which will sting just a bit. It is very important to hold still and let us know if we are causing you discomfort. Next, the doctor will use an X-ray machine (fluoroscope) to guide the needle into the correct spot. An X-ray contrast (dye) may be injected at this point to confirm that the needle has been positioned correctly. Please let the doctor now if you have an allergy to dye. Local anesthetic and steroid are then injected through the needle, and the needle is removed. Your skin will be cleaned and a Band-Aid may be applied. The Band-Aid may be removed the next morning.
What to wear?	Please wear loose, comfortable clothing. Please leave all jewelry and other valuables at home.
How long does it take?	We ask that you arrive 30-45 minutes before the scheduled time of your procedure. The procedure lasts about 15 to 30 minutes, and you will be in the recovery area about 15 minutes.
Can I drive home from the procedure?	No, you must have a ride home, and your driver must remain on site.
Can I eat before?	You may receive conscious sedation. Conscious sedation is medication given through an IV (tube) in your arm. The medication will help you relax, but it will not put you to sleep. If you opt to have conscious sedation, you MUST NOT EAT OR DRINK for <u>4</u> hours before your procedure. You may, however, take your medication with a <u>small sip of water</u> .
Can I take my medications?	Yes, except blood thinners .
What are the possible side effects of steroid medication?	Administration of steroid medication can cause side effects. They include, but not limited to: hyperglycemia (increased blood sugar), sweats, hot/cold flashes, flushing of the face, hypertension (increased blood pressure), osteoporosis, insomnia, mood swings and Cushing's Syndrome.
Risks of this procedure include, but are not limited to:	<ul style="list-style-type: none">* Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol.* <i>Bleeding, infection and/or bruising in the injection area. Increased pain</i>* <i>A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required requiring transportation to the hospital.</i>* A reaction to the local anesthetic or conscious sedation, including but not limited to: allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest requiring transportation to the hospital. .* Temporary numbness or weakness in one or both legs. This is normal.* Nerve damage, paralysis, stroke and even death.

03/15/2015rdsteachingsheetsREVIEWED 3/2016

PLEASE SEE THE BACK FOR YOUR INSTRUCTIONS

Instructions Regarding Your Procedure
Ganglion Impar nerve Block with fluoroscopy

Will I receive a pre-procedure phone call?	A nurse from our office will call you 24-48 hours prior to your procedure. She will ask you a few important medical questions, AND she will confirm your appointment. We must have a verbal confirmation from you 24-48 hours prior to your procedure . Failure to return our call could result in your procedure being rescheduled.
Where can I obtain more procedure information?	Please visit our website www.nyspineandwellness.com . It contains demonstrations of some of the procedures we offer. We also have other helpful information.
Can I drive home from the procedure?	<input type="checkbox"/> You MUST have a ride home, and your driver should remain on site. You must not drive or operate heavy machinery for 24 hours.
Can I eat Breakfast? <i>*the box with a check applies to you</i>	<input type="checkbox"/> Yes, you can eat a light breakfast or light lunch <input type="checkbox"/> No, DO NOT EAT OR DRINK for 4 hours before your procedure. However, you may take your medicine but only with a small sip of water .
Should I stop taking aspirin?	If you take ASPIRIN or products containing ASPIRIN in doses greater than 325 mg/per day, you must stop taking it 7 days prior to your procedure (examples of ASPIRIN-containing products include but are not limited to: EXCEDRIN, FIORINAL, AND ALKA SELTZER, etc).
Can I take the aspirin prescribed for <u>cardiac</u> and or <u>stroke</u> prevention?	<input type="checkbox"/> Yes, you can continue to take your aspirin for cardiac or stroke prevention, not to exceed 325mg per day.
Can I take my blood thinner? <i>*the box with a check applies to you</i>	<input type="checkbox"/> Our provider has advised you on the number of days to hold the blood thinner you take. <input type="checkbox"/> Our provider will have to contact the Doctor who has prescribed the blood thinner for you to learn how long it is safe for you to be off the medicine prior to your procedure. One of our nurses will call you when we have obtained that information.
Can I take my other medications? <i>*the box with a check applies to you</i>	<input type="checkbox"/> Yes, except for blood thinners . <input type="checkbox"/> Yes, only with a small sip of water ,(you must not take your blood thinner)
What time is my procedure?	You will receive a phone call one or two business days prior to your procedure to remind you of the time you should arrive for your procedure (usually 30-40 minutes prior to the scheduled time) and the location. PLEASE BE ON TIME . <i>*some patients may receive a call to confirm insurance information.</i>
How long will it take?	We ask that you plan <u>1½ -3 hours</u> from the time you arrive at the center to the time you are discharged. Please understand that circumstances sometimes arise that are out of our control, which cause a longer delay. Please plan accordingly.
What do I bring?	You must bring the following <u>each time</u> you have a procedure: 1. Your current insurance information 2. A written list of all your current medications including the strength and frequency you take them. Please include over the counter medication, vitamins and herbal supplements.
What shall I wear?	Please wear comfortable loose fitting clothing.
Can I wear jewelry?	No. Please leave all jewelry and other valuables at home. We are unable to store them safely. Any body- piercing jewelry must be completely removed, as this may interfere with the x-ray machine.
What do I do if I am ill or have to cancel?	Please call the office 24 hours in advance if you are unable to keep your appointment. (315) 552-6700. We do charge patients for no shows.