

Splanchnic Block

11/14 ! !49	A sulpushuis blook is an injection administrated into the suine. A flyanessame	
What is it?	A splanchnic block is an injection administered into the spine. A fluoroscope	
	(x-ray machine) assists the physician in locating the nerve roots that are	
	inflamed and contributing to your pain.	
Why is it done?	This procedure is performed to diagnose or treat pain that originates in the chest	
	and upper abdomen (pancreas, liver, stomach or upper intestine). Nerves that	
	serve the autonomous (involuntary) nervous system and abdominal organs are	
	located in front of the spinal column. Sometimes a series of blocks is required	
	to achieve maximum benefit.	
How is it done?	When you are in the procedure room, you will be asked to lie face down on a	
	cushioned x-ray table. A small needle is used to inject a local anesthetic into the	
	areas around the injection site to numb the skin. Next, a long thin needle is	
	placed in the side of the mid to upper back and advanced to the front of the	
	spinal column; placement is confirmed with the fluoroscope and injection of	
	contrast dye. After placement of one or two needles a solution of local	
	anesthetic is injected. The medication will travel through the spinal column to	
	the sympathetic nerves in the chest and abdomen. The needles are removed and	
	small band aids are applied.	
Is there any	You may receive conscious sedation. Conscious sedation is medication given	
preparation?	through an intravenous (IV) catheter (tube) in a vein in your arm. The	
	medication will help you relax, but it will not put you to sleep. If you opt to	
	have conscious sedation, you MUST NOT EAT OR DRINK for 4 hours	
	before your procedure. However, you may take your medication with a	
	small sip of water. You must have someone available to drive you home	
	after the procedure.	
What to wear?	Please wear loose, comfortable clothing and leave all jewelry/other valuables at	
	home.	
How long does	We ask that you arrive 30-45 minutes before the scheduled time of your	
it take?	procedure. The procedure lasts about 15 - 30 minutes, and you will be in the	
	1 415 : 4	
	recovery area about 15 minutes.	
	· ·	
Risks of this	* Pain in the area where the needle(s) was inserted. The pain can last for two	
Risks of this procedure	* Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain	
Risks of this procedure include, but	* Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol.	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further 	
Risks of this procedure include, but	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. A reaction to the local anesthetic or conscious sedation, including but not 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. A reaction to the local anesthetic or conscious sedation, including but not limited to: Temporary numbness or weakness in one or both legs. This is 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. A reaction to the local anesthetic or conscious sedation, including but not limited to: Temporary numbness or weakness in one or both legs. This is normal. You should have assistance with walking and should not drive for 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. A reaction to the local anesthetic or conscious sedation, including but not limited to: Temporary numbness or weakness in one or both legs. This is 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. A reaction to the local anesthetic or conscious sedation, including but not limited to: Temporary numbness or weakness in one or both legs. This is normal. You should have assistance with walking and should not drive for 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. A reaction to the local anesthetic or conscious sedation, including but not limited to: Temporary numbness or weakness in one or both legs. This is normal. You should have assistance with walking and should not drive for 24 hours. Bleeding, bruising, or infection in the injection area. 	
Risks of this procedure include, but are not limited	 Pain in the area where the needle(s) was inserted. The pain can last for two to three days and can be treated by using ice and mild analgesics (pain medication) such as Motrin, Naprosyn or Tylenol. A reaction to the contrast dye. These reactions usually do not require further treatment. However, a reaction to the contrast dye may result in sneezing, hives, swelling of the face, and throat, respiratory difficulties and shock. Medication and/or respiratory assistance may be required. Allergic reaction to the anesthetic medication, loss of consciousness, depressed breathing and cardiac arrest. A reaction to the local anesthetic or conscious sedation, including but not limited to: Temporary numbness or weakness in one or both legs. This is normal. You should have assistance with walking and should not drive for 24 hours. Bleeding, bruising, or infection in the injection area. 	

09-12-2015;06-10-2007 rdswteachingsheet REVIEWED 3/2016 REVISED 07-07-2016 reviewed with changes 09-29-2020

Instructions Regarding Your Procedure Splanchnic Block

Will I receive a pre-procedure phone call?	A nurse from our office will call you 24-48 hours prior to your procedure. She will ask you a few important medical questions, <u>AND</u> she will confirm your appointment. We must have <u>a verbal confirmation from you 24-48 hours prior to your procedure.</u> Failure to return our call could result in your procedure being rescheduled. The number to call us back is: (315) 703-3480
Where can I obtain more	Please visit our website www.nyspineandwellness.com. It contains demonstrations of some of
procedure information?	the procedures we offer. We also have other helpful information.
Can I drive home from the	·
procedure?	You MUST have a ride home, and your driver should remain on site. You must not drive or operate heavy machinery for 24 hours.
Can I eat Breakfast?	☐ No, DO NOT EAT OR DRINK for 4 hours before your procedure. However, you may take
	your medicine but only with a <u>small sip of water</u> .
Should I stop taking aspirin?	If you take ASPIRIN or products containing ASPIRIN for <u>PAIN or HEADACHES</u> , you <u>MUST STOP</u> 7 days prior to procedure. Products include but are not limited to: EXCEDRIN, FIORINAL, AND ALKA SELTZER, etc).
Can I take the aspirin	
prescribed for cardiac and or	☐ Yes, however you need to limit the daily dose to 81 mgs per day for the 7 days prior to the
stroke prevention?	day of the procedure.
Can I take my blood thinner? *the box with a check applies to you	☐ Our provider has advised you on the number of days to hold the blood thinner you take. ☐ Our provider will have to contact the Doctor who has prescribed the blood thinner for you
	to learn how long it is safe for you to be off the medicine prior to your procedure. One of our nurses will call you when we have obtained that information.
Can I take my other	Yes, except for blood thinners .
medications?	Yes, only with a small sip of water, (you must not take your blood thinner)
*the box with a check applies to you	
What time is my procedure?	You will receive a phone call one or two business days prior to your procedure to remind you
	of the time you should arrive for your procedure (usually 30-40 minutes prior to the scheduled
	time) and the location. PLEASE BE ON TIME.
	*some patients may receive a call to confirm insurance information.
How long will it take?	We ask that you plan 1½ -3 hours from the time you arrive at the center to the time you are
	discharged. Please understand that circumstances sometimes arise that are out of our control,
	which cause a longer delay. Please plan accordingly.
What do I bring?	You <i>must</i> bring the following <u>each time</u> you have a procedure:
	1. Your current insurance information
	2. A written list of all your current medications including the strength and frequency you take
	them. Please include over the counter medication, vitamins and herbal supplements.
What shall I wear?	Please wear comfortable loose fitting clothing.
Can I wear jewelry?	No. Please leave all jewelry and other valuables at home. We are unable to store them safely.
can't wear jeweny:	Any body- piercing jewelry must be completely removed, as this may interfere with the x-ray
	machine.
What do I do if I am ill or	Please call the office 24 hours in advance if you are unable to keep your appointment.
have to cancel?	(315) 552-6700. We do charge patients for no shows.
nave to cancer:	(313) 332 0700. We do charge patients for no shows.